

Acacia mellifera

(Vahl) Benth.

Fabaceae - Mimosoideae

LOCAL NAMES

Afrikaans (swarthook,swartaak); Arabic (kitr,kedad,kitir); English (wait-a-bit thorn,black thorn,hook thorn); Ndebele (umngaga,mukotokwa,muguhungu,katogwa); Somali (bilel,laner,lanen); Swahili (kikwata); Tigrigna (tselim kenteb); Tongan (mupandabutolo); Tswana (wynruit,mongana,hakiesdoring,haakdoring,blouhaak)

BOTANIC DESCRIPTION

Acacia mellifera is a low, branched tree with a more or less spherical crown. Black bark on stem becomes ash-grey to light brown on the branches, bearing small, short, sharply hooked spines in pairs. It has a shallow but extensive root system radiating from the crown, allowing the plant to exploit soil moisture and nutrients from a large volume of soil. The roots rarely penetrate more than 1 m.

Leaves characterized by 2 pairs of pinnulae, each with a single pair of leaflets. Leaflets elliptic 0.6-2 cm long and 0.6-1.2 cm wide, glabrous and highly coloured beneath.

Flowers sweetly scented, especially at night, in elongated spikes, cream to white in spiciform racemes, up to 3.5 cm long; pedicels 0.5-1.5 mm long; calyx up to 1 mm long; corolla 2.5-3.5 mm long.

The papery pods with 2-3-seeds are reticulate, flat, elongated, 2.5-5.5 cm long, 6 cm wide, hemmed, sometimes more or less narrowed between the seeds.

The generic name 'acacia' comes from the Greek word 'akis', meaning point or barb. The specific name means 'honey-bearing'.

BIOLOGY

Flowering and fruiting start 3 years after planting and generally occur twice a year. Flowers are borne on shoots produced the previous year, that is, old wood. The flowers open and shed before the leaves appear. In southern Africa, flowering occurs from September to November and fruiting from January to April.



Tree habit: *A. mellifera* ssp. *mellifera* in the Rift valley, Kenya. (Chris Fagg)



Tree habit: Multistemmed habit of *A. mellifera* ssp. *mellifera* close to the Angolan border in Namibia. (Chris Fagg)



Acacia mellifera foliage (Chris Fagg)

ECOLOGY

A. mellifera is a commonly occurring shrub on rangelands throughout the savannah in western, eastern and southern Africa. The terrain preference is rocky hillsides with rainfall along seasonal watercourses, mixed with other trees. If left unattended, especially if grazing is heavy and no fires check its spread, it may form dense, impenetrable thickets, 2-3 m high and sometimes hundreds of metres across, slowly taking over good grazing land. This species is drought-tolerant.

BIOPHYSICAL LIMITS


Altitude: 0-1500 m, Mean annual rainfall: 250-700 mm

Soil type: *A. mellifera* is normally found on hard-surfaced, sandy soils and rocky hillsides. It grows well in black cotton soils but prefers loamy soils.

DOCUMENTED SPECIES DISTRIBUTION

Native: Angola, Botswana, Chad, Eritrea, Ethiopia, Kenya, Namibia, Oman, Saudi Arabia, Somalia, Sudan, Tanzania, Uganda, Yemen, Republic of, Zambia, Zimbabwe

Exotic: Mozambique, South Africa



The map above shows countries where the species has been planted. It does neither suggest that the species can be planted in every ecological zone within that country, nor that the species can not be planted in other countries than those depicted. Since some tree species are invasive, you need to follow biosafety procedures that apply to your planting site.

PRODUCTS

Food: Gum collected from injured stems is edible and relished by children, animals and birds.

Fodder: Camels and goats browse the leaves, which are rich in protein, taking them from the shrubs or from the ground.

Apiculture: The cream/white flowers produce excellent quality honey ('mellifera' = producing honey). Bees forage in the late morning to mid afternoon when hot and dry. The honey is water colour and granulates slowly.

Fuel: The wood is used for fuel and charcoal.

Timber: The wood is taken for building huts and the branches for fencing.

Poison: The poison with which Bushmen tip their arrows is often made from a powdered grub mixed with the sap of *A. mellifera*.

Medicine: The bark decoction is used for stomach-ache, sterility, pneumonia, malaria and syphilis. In Botswana, a decoction of the roots is a medicine for stomach pain.

Other products: In Sudan, baskets made of the roots serve for collecting gum arabic. The bast fibres are used for wickerwork.

SERVICES

Shade or shelter: When pruned regularly, *A. mellifera* in arid areas provides animals with shade from the hot sun.

Boundary or barrier or support: Makes a good live fence and hedge.

Intercropping: *A. mellifera* has an aggressive root system, limiting its use in farms with crops.

TREE MANAGEMENT

Young trees are subject to heavy browsing by stock and game and must be protected for the first two seasons. *A. mellifera* has a moderate growth rate of up to 500 mm/year. It does not coppice well.

GERMPLASM MANAGEMENT

Seed storage behaviour is orthodox; viability can be maintained for several years in hermetic storage at 10 deg. C with 4.5-9% mc. There are approximately 20 000 seeds/kg.

FURTHER READING

- Albrecht J. ed. 1993. Tree seed hand book of Kenya. GTZ Forestry Seed Center Muguga, Nairobi, Kenya.
- Beentje HJ. 1994. Kenya trees, shrubs and lianas. National Museums of Kenya.
- Bein E. 1996. Useful trees and shrubs in Eritrea. Regional Soil Conservation Unit (RSCU), Nairobi, Kenya.
- CABI. 2000. Global Forestry Compendium. CD-ROM. CABI
- Coates-Palgrave K. 1988. Trees of southern Africa. C.S. Struik Publishers Cape Town.
- Dale IR, Greenway PJ. 1961. Kenya trees and shrubs. Buchanan's Kenya Estates Ltd.
- Eggeling. 1940. Indigenous trees of Uganda. Govt. of Uganda.
- El Amin HM. 1973. Sudan acacias. Forest Research Institute Publishing Section Information Department.
- FAO, UNEP. 1983. Notes on trees and shrubs in arid and semi-arid regions. EMASAR phase II. FAO, Rome.
- Hassan AS, Styles BT. 1990. A conspectus of Somali Acacias. Chatham: Natural Resource Institute, Oxford Forestry Institute. Somali Forestry Series No. 4.
- Hines DA, Eckman K. 1993. Indigenous multipurpose trees for Tanzania: uses and economic benefits to the people. Cultural survival Canada and Development Services Foundation of Tanzania.
- Hong TD, Linington S, Ellis RH. 1996. Seed storage behaviour: a compendium. Handbooks for Genebanks: No. 4. IPGRI.
- ICRAF. 1992. A selection of useful trees and shrubs for Kenya: Notes on their identification, propagation and management for use by farming and pastoral communities. ICRAF.
- Katende AB et al. 1995. Useful trees and shrubs for Uganda. Identification, Propagation and Management for Agricultural and Pastoral Communities. Regional Soil Conservation Unit (RSCU), Swedish International Development Authority (SIDA).
- Mbuya LP et al. 1994. Useful trees and shrubs for Tanzania: Identification, Propagation and Management for Agricultural and Pastoral Communities. Regional Soil Conservation Unit (RSCU), Swedish International Development Authority (SIDA).
- Milton SJ. 1986. Phenology of seven Acacia species in Southern Africa. African Journal of Wildlife Research. 17(1): 1-6.
- Noad T, Birnie A. 1989. Trees of Kenya. General Printers, Nairobi.
- Palmer E, Pitman N. 1972. Trees of Southern Africa Vol. 2. A.A. Balkema Cape Town.
- Sahni KC. 1968. Important trees of the northern Sudan. United Nations and FAO.
- Tietema T, Merkesdal E and Schroten J. 1992. Seed germination of indigenous trees in Botswana. Acts Press.
- Timberlake J, Fagg C and Barnes R. 1999. Field guide to the Acacias of Zimbabwe. CBC Publishings, Zimbabwe.
- Timberlake J. 1980. Handbook of Botswana Acacias. Ministry of Agriculture, Botswana.
- Venter F, Venter J-A. 1996. Making the most of Indigenous trees. Briza Publications.
- Vogt K. 1995. A field guide to the identification, propagation and uses of common trees and shrubs of dryland Sudan. SOS Sahel International (UK).

SUGGESTED CITATION

Orwa C, A Mutua, Kindt R, Jamnadass R, S Anthony. 2009 Agroforestry Database: a tree reference and selection guide version 4.0 (<http://www.worldagroforestry.org/sites/treedbs/treedatabases.asp>)