

## Ficus sycomorus

L.

Moraceae

### LOCAL NAMES

Afrikaans (sycomorusvy,rivierwildevy); Amharic (shola,bamba); Arabic (subula); Bemba (mkunyu); English (wild fig, strangler-fig, Sycamore, sycamore fig, bush fig, common cluster fig); French (figuier sycamore, Sykomore); Lozi (mukuyu, katema); Luganda (mukunyu); Lunda (mukuyu); Nyanja (mkuyu); Somali (barda); Spanish (sicomoro); Swahili (mukuyu, mkuyu, chivuzi); Tigrigna (saghla, sagla, shegla); Tongan (mukuyu); Zulu (umKhiwane)

### BOTANIC DESCRIPTION

*Ficus sycomorus* is a large, semi-deciduous spreading savannah tree, up to 21 (max. 46) m, occasionally buttressed. Bark on young stems pale green with a soft powdery covering; on older stems, grey-green, fairly smooth, with scattered grey scales and pale brown patches where scales have fallen off. Slash pale pink with heavy latex flow.

Leaves broadly (ob)ovate or elliptic, base (sub)cordate, apex rounded or obtuse, margin entire or slightly repand -dentate, 2.5-13 (max. 21) x 2-10 (max. 16) cm, scabrous above, petiole 1-5 cm, 5-7 pairs of yellow lateral veins, lowest pair originating at the leaf base.

Flowers, unisexual, cyclic and greenish.

Figs in leaf axils or on up to 10 cm leafless branches on old wood, solitary or paired, globose or (ob)ovoid, yellow-red to reddish-purple when ripe, up to 3.5 x 5 cm, pubescent or almost glabrous. Seeds, numerous, round and very tiny.

*Ficus* is the Latin for fig, derived from the Persian 'fica'. In Greek 'syka' means fig. The species name comes from the Greek 'sykamorea' (sycamore), used in the Gospel according to St. Luke; it was such a tree that Jesus cursed because it was barren. But the word 'sykomorom' had been used to denote the fruit a century before Christ. It has since been applied as a popular name to many sorts of tree, including *Acer pseudoplatanus* and *Platanus occidentalis*.

### BIOLOGY

In southern Africa, flowering and fruiting occur throughout the year, with a peak from July to December. Small wasps (*Ceratosolen arabicus*), which develop in some of the flowers and live symbiotically inside the syconium, pollinate the unisexual flowers. Bats achieve seed dispersal.



(Joris de Wolf, Patrick Van Damme, Diego Van Meersschaut)



*Ficus sycomorus* bark (Joris de Wolf, Patrick Van Damme, Diego Van Meersschaut)



(Joris de Wolf, Patrick Van Damme, Diego Van Meersschaut)

**ECOLOGY**

A common savannah tree that grows in high water table areas. Often found along watercourses such as streams and rivers, swamps and waterholes. The sycamore fig is sensitive to frost but can withstand some cold. It is found in afro-montane rain forests and undifferentiated afro-montane forests, especially along edges and in clearings, riverine forests, riparian woodland, secondary evergreen bushland; left as single trees in farmland and occasionally seen as single trees on rocky outcrops.

**BIOPHYSICAL LIMITS**

Altitude: 0-2000 m, Mean annual temperature: 0-40 deg. C, Mean annual rainfall: 500-1800 (max. 2200) mm

Soil type: Prefers deep, well-drained loam to clay soil rich in nutrients. Sandy soils with a shallow groundwater level may also be suitable.

**DOCUMENTED SPECIES DISTRIBUTION**

Native: Angola, Benin, Botswana, Burundi, Cameroon, Congo, Cote d'Ivoire, Democratic Republic of Congo, Djibouti, Egypt, Eritrea, Ethiopia, Gambia, Ghana, Guinea, Israel, Kenya, Lesotho, Madagascar, Mozambique, Namibia, Nigeria, Rwanda, Saudi Arabia, Senegal, Somalia, South Africa, Sudan, Swaziland, Syrian Arab Republic, Tanzania, Uganda, Zambia, Zimbabwe

Exotic:



The map above shows countries where the species has been planted. It does neither suggest that the species can be planted in every ecological zone within that country, nor that the species can not be planted in other countries than those depicted. Since some tree species are invasive, you need to follow biosafety procedures that apply to your planting site.

**PRODUCTS**

**Food:** Mature fruits are eaten fresh, stewed, or dried and stored for later use. Fruit can also be used for the preparation of an alcoholic beverage. Leaves are used in soups and groundnut dishes. The bark is chewed together with kola nut. In Ghana, the wood ash is commonly used as a salt substitute.

**Fodder:** Leaves are a much-sought fodder with fairly high nutritive value (9% crude protein and 7 mJ/kg net energy dry matter); they are valuable fodder in overstocked semi-arid areas where the trees occur naturally. Fruits are eaten by livestock, wild animals and birds.

**Fuel:** Can be used as firewood and for making charcoal; various peoples throughout Africa use a piece of dry wood from this tree as the base block when starting a fire by the friction method.

**Fibre:** Inner part of the root used as weaving fibre, and a strong rope can be made from the inner bark.

**Timber:** The wood is creamy brown, has a fairly uniform structure, is very light (air-dry 510 kg/m<sup>3</sup>), soft to moderately hard, tough, strong, easy to work, finishes smoothly and holds nails firmly. It is not very durable and is easily attacked by termites. Mainly used for making mortars and pestles, drums, stools, doors, beehives, dugout canoes, carvings and for house building.

**Medicine:** The bark is used for the treatment of scrofula, coughs, and throat and chest diseases. The milky latex is used for treatment of dysentery and chest diseases, or is applied to inflamed areas, while ringworm is treated with the bark and milky latex. Leaves are said to be effective against jaundice and as an antidote for snakebite, while the roots have laxative and anthelmintic properties.

**SERVICES**

**Erosion control:** Wild fig can be used for sand-dune fixation and riverbank stabilization.

**Shade or shelter:** The tree gives useful shade and is common at marketplaces, where people gather under it for many social functions.

**Soil improver:** Shed leaves form a valuable litter improving the nutrient status, infiltration rate and water-holding capacity of the soil.

**Ornamental:** An important tree planted for ornamental purposes near temples, roadsides, wells and community places such market centres in rural areas. The yellowish bark shows at an early stage, contrasting well with the green leaves. A popular species to grow as a bonsai.

**Intercropping:** Usually intercropped with bananas as an understory.

**Other services:** The sycamore fig is widely valued for spiritual and sacred purposes. Such use can be traced back to ancient Egypt, and it is often mentioned in the Holy Bible.

**TREE MANAGEMENT**

The sycamore fig is large, spreading and very shady, and therefore requires considerable spacing. It tolerates lopping. Continuous and adequate water supply is necessary for high yields, but rainfall during fruiting may cause fruits to split and favour growth of fungal rots. The best site for trees is next to drainage lines, streams, rivers, springs or dams. A tree can bear several crops of fruit a year. The invasive root system should be borne in mind when positioning trees. Wild fig does well in a container if pruned once or twice a year. Growth rate is fairly fast at 1-1.5 m/year in frost-free areas.

**GERMPLASM MANAGEMENT**

Viable seed is difficult to obtain.

**PESTS AND DISEASES**

The larvae of several Lepidoptera and Coleoptera (long-horn beetles) make tunnels in the branches and sometimes the trunk. Many leaf-eating beetles (Coleoptera) and caterpillars (Lepidoptera) damage foliage. Scale insects and mealybugs (Homoptera), which suck the sap, may attack fruits, branches and leaves. The larvae of fruit flies of *Dacus* spp. and *Ceratitis* spp. (Diptera) feed on the pulp of fruits, resulting in rotting and premature drops. Other pests include *Drosophila* spp., *Acrina* spp., *Aceria ficus* and a sawfly (Hymenoptera).

At least 30 species of fungus attack the genus *Ficus*. These include root rots, branch wilt and canker, leaf rusts, branch and foliage blights, fruit surface mould and spot rot, internal fruit rot, mould and smut and fruit souring. Most are of minor importance and can be controlled by chemical sprays. The tree is susceptible to nematodes, hence should not be planted in infested soils.

**FURTHER READING**

- Abbiw D. 1990. Useful plants of Ghana. Intermediate Technology Publications and the Royal Botanical Gardens, Kew.
- Beentje HJ. 1994. Kenya trees, shrubs and lianas. National Museums of Kenya.
- Bein E. 1996. Useful trees and shrubs in Eritrea. Regional Soil Conservation Unit (RSCU), Nairobi, Kenya.
- Bekele-Tesemma A, Birnie A, Tengnas B. 1993. Useful trees and shrubs for Ethiopia. Regional Soil Conservation Unit (RSCU), Swedish International Development Authority (SIDA).
- Coates-Palgrave K. 1988. Trees of southern Africa. C.S. Struik Publishers Cape Town.
- Crane E, Walker P. 1984. Pollination directory for world crops. International Bee Research Association, London, UK.
- Dale IR, Greenway PJ. 1961. Kenya trees and shrubs. Buchanan's Kenya Estates Ltd.
- Eggeling. 1940. Indigenous trees of Uganda. Govt. of Uganda.
- Friis I. 1992. Forests and forest trees of northeast tropical Africa. Her Majesty's Stationery Office, London.
- Glenn WH. 1988. Insect enemies of trees. Logos Press. New Delhi. India.
- Hines DA, Eckman K. 1993. Indigenous multipurpose trees for Tanzania: uses and economic benefits to the people. Cultural survival Canada and Development Services Foundation of Tanzania.
- ICRAF. 1992. A selection of useful trees and shrubs for Kenya: Notes on their identification, propagation and management for use by farming and pastoral communities. ICRAF.
- International Board for Plant Genetic Resources (IBPGR). 1986. Genetic Resources of Tropical and sub-Tropical Fruits and Nuts.
- Katende AB et al. 1995. Useful trees and shrubs for Uganda. Identification, Propagation and Management for Agricultural and Pastoral Communities. Regional Soil Conservation Unit (RSCU), Swedish International Development Authority (SIDA).
- Lanzara P. and Pizzetti M. 1978. Simon & Schuster's Guide to Trees. New York: Simon and Schuster
- Mariella P, Lanzara P. 1977. Guide to trees. Simon and Schuster.
- Mbuya LP et al. 1994. Useful trees and shrubs for Tanzania: Identification, Propagation and Management for Agricultural and Pastoral Communities. Regional Soil Conservation Unit (RSCU), Swedish International Development Authority (SIDA).
- Noad T, Birnie A. 1989. Trees of Kenya. General Printers, Nairobi.
- Peter G von Carlowitz. 1991. Multipurpose Trees and Shrubs-Sources of Seeds and Inoculants. ICRAF. Nairobi, Kenya
- Record SJ. 1972. Timbers of the New World. Yale University Press.
- Storrs AEG. 1995. Know your trees: some common trees found in Zambia. Regional Soil Conservation Unit (RSCU).
- Szolnoki TW. 1985. Food and fruit trees of Gambia. Hamburg. Federal Republic of Germany.
- Venter F, Venter J-A. 1996. Making the most of Indigenous trees. Briza Publications.
- von Maydell HJ. 1986. Trees and shrubs of the Sahel - their characteristics and uses. GTZ 6MBH, Eschborn.

**SUGGESTED CITATION**

Orwa C, A Mutua, Kindt R, Jamnadass R, S Anthony. 2009 Agroforestry Database: a tree reference and selection guide version 4.0 (<http://www.worldagroforestry.org/sites/treedbs/treedatabases.asp>)