

LOCAL NAMES

English (male funtum,Lagos rubber,bush rubber,bastard wild rubber)

BOTANIC DESCRIPTION

Funtumia africana is a tropical tree up to 30 m tall (usually shorter) with a straight, cylindrical trunk and a narrow tree crown. Bark brown to dark in colour, thin and slightly fissured becoming granular on old trees. Slash orange exuding latex copiously.

Leaves elliptic or ovate, base round or cuneate, apex acuminate 20 x 9 cm, with approximately 8-14 main lateral veins on each side, leaf margins wavy. Axils on the main lateral veins not pitted.

Flowers yellow-white, fragrant in dense cymes. Corolla tube 6-10 mm, lobes 5-7mm.

Fruit grey-brown, fusiform, with an acute or acuminate apex, up to 30 cm long, with hairy wind borne seeds.

Funtumia africana and *F. elastica* are very similar morphologically and distribution wise, when *F. elastica* latex is rubbed between the fingers the latex coagulates into balls, whereas *F. africana* does not. The generic epithet is derived from 'funtum', a local Ghanaian (Akan dialect) name of the plant. The specific epithet means 'of Africa'.

BIOLOGY

Flowers insect pollinated.

ECOLOGY

A forest tree commonly composing the second storey in rainforests, *F. africana* is commonly associated with *Elaeis guineensis*, *Kolobopetalum chevalieri*, *Dioscorea preusii*, *Cnestis hirsutus*, *Tabernaemontana crassa* and *Alstonia congensis*, this association is the *Tabernaemontano-Elaeetum guineensis* typicum. Bastard wild rubber is also common in forest regrowth.

DOCUMENTED SPECIES DISTRIBUTION

Native: Angola, Cameroon, Cote d'Ivoire, Democratic Republic of Congo, Gabon, Kenya, Liberia, Mozambique, Nigeria, Sierra Leone, Tanzania, Togo, Uganda

Exotic:



The map above shows countries where the species has been planted. It does neither suggest that the species can be planted in every ecological zone within that country, nor that the species can not be planted in other countries than those depicted. Since some tree species are invasive, you need to follow biosafety procedures that apply to your planting site.

PRODUCTS

Apiculture: The fragrant yellow-white flowers are visited by bees for nectar and pollen.

Fuel: Branches and trunk can be used as fuelwood.

Timber: *F. africana* wood is white, even textured with a weight of 64 kg/m³. air dry. It is used for cheap joinery, furniture and matchstick manufacture. The wood is also reportedly used for carving stools, doors and miscellaneous household requirements.

Gum or resin: In Kenya, *F. africana* latex is used as birdlime.

Latex or rubber: *F. africana* latex polyisoprenes are useless as a source of rubber, Use of bastard wild rubber latex as an adulterant, leads to imperfect coagulation and consequently a valueless rubber.

Medicine: In Africa this species is used to treat urinary incontinence and burns. The leaf and bark are used as enema. The principle alkaloids of *F. africana*, funtumine and funtumidine are hypotensive.

Poison: The latex is a weak ingredient and adherent for poisoned arrows.

Other products: Conanine-type alkaloids were isolated from the stem bark (collected in Nigeria), some are identical with holonamine isolated from the stem bark of *Holarrhena antidysenterica*. The extracts induced a dose-related bradycardia in rats and, in vitro, in guinea pig auricles. The seed floss is used to stuff pillows in Ghana.

SERVICES

Erosion control: The root system of *F. africana* protects moist forest soil from erosion.

Shade or shelter: The tree casts a heavy, cool shade.

Soil improver: Leaf litter improves soil fertility of surrounding soil.

FURTHER READING

Abbas B, El-Tayeb AE, Sulleiman YR. 1992. *Calotropis procera*: feed potential for arid zones. *Veterinary-Record*. 131(6):132.

Beentje HJ. 1994. Kenya trees, shrubs and lianas. National Museums of Kenya.

Burkill HM. 1994. Useful plants of West Tropical Africa. Vol. 2. Families E-I. Royal Botanical Gardens, Kew.

Eggeling. 1940. Indigenous trees of Uganda. Govt. of Uganda.

Hamilton A.C. 1981. A field guide to Uganda forest trees.

Masens BYD et al. 1996. The association of *Elaeis guineensis* and *Tabernaemontana crassa* in the region of Kikwit (Zaire). *Fragmenta Floristica et Geobotanica*. 41 (1): 295-313.

Omino EA and Kokwaro JO. 1993. Ethnobotany of Apocynaceae species in Kenya. *Journal of Ethnopharmacology*. 40(3): 167-180.

Wagner H et al. 1987. The steroid alkaloids of *Funtumia africana*. *Planta Medica*. 53(5): 444-449.

SUGGESTED CITATION

Orwa C, A Mutua, Kindt R, Jamnadass R, S Anthony. 2009 *Agroforestry Database: a tree reference and selection guide version 4.0* (<http://www.worldagroforestry.org/sites/treedbs/treedatabases.asp>)