

**LOCAL NAMES**

Amharic (kosso); English (kousso, hagenia, cusso, brayera, African redwood); Swahili (mlozilozi, mdobore)

**BOTANIC DESCRIPTION**

*Hagenia abyssinica* is a slender tree up to 20 m tall, with a short trunk and thick branches; branchlets covered in silky brown hairs and ringed with leaf scars. Bark thick, brown or reddish-brown and readily peeling. No thorns or buttresses.

Leaves compound, 40 cm long, in terminal tufts; leaflets pale or bright green above, with silvery hairs below, reddish and sticky when young, 3-6 pairs plus a terminal leaflet, each about 10 cm long; margin finely toothed and fringed with long hairs; leaf stalks 12 cm long, with expanded wings formed from the stipules, densely hairy on the underside.

Flowers in handsome multibranched, terminal, drooping panicles up to 60 cm long and 30 cm wide, polygamo-dioecious, female heads pinkish-red, clearly veined, bulkier than the more feathery orange-buff to white male heads.

Fruit small, dry, winged, asymmetric, single seeded, brown syncarp with a single more or less ovoid carpel and fragile pericarp.

*Hagenia* is a monospecific genus and is most closely related to the monospecific genus *Leucosidea*. The specific name means 'from Ethiopia'.

**BIOLOGY**

Trees have either male or female flowers. Flowering and seeding can be observed throughout the year with a break in the months with the coldest temperatures.

**ECOLOGY**

First described in Ethiopia and also found in East Africa, this tree is often dominant in the woodland zone just above the mountain bamboo. In Kenya, the species can be found on the slopes of Mt. Kenya, Mt. Elgon, the Aberdares, Cherangani Hills and the slopes of other high hills or mountains. It occurs in undifferentiated afro-montane forest (mixed Podocarpus forest, Juniperus-Podocarpus forest) and dry single dominant afro-montane forest (Juniperus forest or forest dominated by Hagenia), especially along the upper limit; often associated with Schefflera abyssinica, S. volkensii, Galiniera saxifraga, Rapanea melanophloës and with the mountain bamboo, Arundinaria alpina; at lower altitudes often at forest margins. This is a very clear example of an afro-montane endemic.

**BIOPHYSICAL LIMITS**


Altitude: 2000-3000 m

Mean annual rainfall: 1000-1500 mm

**DOCUMENTED SPECIES DISTRIBUTION**

Native: Burundi, Democratic Republic of Congo, Ethiopia, Kenya, Malawi, Rwanda, Sudan, Tanzania, Uganda, Zambia

Exotic: India



The map above shows countries where the species has been planted. It does neither suggest that the species can be planted in every ecological zone within that country, nor that the species can not be planted in other countries than those depicted. Since some tree species are invasive, you need to follow biosafety procedures that apply to your planting site.

**PRODUCTS**

Fuel: *H. abyssinica* is a good source of firewood and charcoal.

Timber: Wood is dark red, medium soft but not durable; it is used for furniture, poles, flooring, carving and cabinet making.

Medicine: The roots are cooked with meat and the soup drunk for general illness and malaria, while the dried and pounded female inflorescence is used as an anthelmintic (especially for tapeworm). Bark may be pounded, added to cold water and the liquid drunk as a remedy for diarrhoea and stomach-ache. Generally, this is a strong medicine that must not be taken in large quantities; it is sometimes taken as an abortifacient.

**SERVICES**

Erosion control: Trees are employed in soil-conservation activities.

Shade or shelter: The fire-resistant species can be used as a firebreak.

Soil improver: *Hagenia* constantly sheds leaves, providing mulch and green manure.

Ornamental: The attractive tree is suitable for planting in amenity areas.

Intercropping: In an experiment in Makete District Tanzania, grain yield of wheat was measured on eight different farms for a two-year period. The threshed wheat yield under a pruned *H. abyssinica* canopy averaged 810 kg/ha compared to 600kg/ha on parts of the field located at least 30 m from a tree.

**TREE MANAGEMENT**

In Tanzania, trees are planted or wildings are allowed to grow, interspersed throughout a field at a density of about 10 trees/ha. To reduce light competition with crop and foster marketable bole development, the trees are pruned once each year in November, about two months before planting crops at the onset of the rainy season. Typically, annual prunings of mature trees remove 40-50% of the canopy height. Pollarding is also a suitable silvicultural practice.

**GERMPLASM MANAGEMENT**

Seed storage behaviour is orthodox; hermetic air-dry storage at cool temperatures is recommended. On average, there are 400 000-500 000 seeds/kg.

**PESTS AND DISEASES**

Wood is susceptible to attack by borers.

**FURTHER READING**

Abbas B, El-Tayeb AE, Sulleiman YR. 1992. *Calotropis procera*: feed potential for arid zones. *Veterinary-Record*. 131(6):132.

Albrecht J. ed. 1993. *Tree seed hand book of Kenya*. GTZ Forestry Seed Center Muguga, Nairobi, Kenya.

Anon. 1986. *The useful plants of India*. Publications & Information Directorate, CSIR, New Delhi, India.

Beentje HJ. 1994. *Kenya trees, shrubs and lianas*. National Museums of Kenya.

Bekele-Tesemma A, Birnie A, Tengnas B. 1993. *Useful trees and shrubs for Ethiopia*. Regional Soil Conservation Unit (RSCU), Swedish International Development Authority (SIDA).

Dale IR, Greenway PJ. 1961. *Kenya trees and shrubs*. Buchanan's Kenya Estates Ltd.

Friis I. 1992. *Forests and forest trees of northeast tropical Africa*. Her Majesty's Stationery Office, London.

Hamilton A.C. 1981. *A field guide to Uganda forest trees*.

Hines DA, Eckman K. 1993. *Indigenous multipurpose trees for Tanzania: uses and economic benefits to the people*. Cultural survival Canada and Development Services Foundation of Tanzania.

Hong TD, Linington S, Ellis RH. 1996. *Seed storage behaviour: a compendium*. Handbooks for Genebanks: No. 4. IPGRI.

ICRAF. 1992. *A selection of useful trees and shrubs for Kenya: Notes on their identification, propagation and management for use by farming and pastoral communities*. ICRAF.

Katende AB et al. 1995. *Useful trees and shrubs for Uganda. Identification, Propagation and Management for Agricultural and Pastoral Communities*. Regional Soil Conservation Unit (RSCU), Swedish International Development Authority (SIDA).

Kokwaro JO. 1976. *Medicinal plants of East Africa*. East African Literature Bureau.

Mbuya LP et al. 1994. *Useful trees and shrubs for Tanzania: Identification, Propagation and Management for Agricultural and Pastoral Communities*. Regional Soil Conservation Unit (RSCU), Swedish International Development Authority (SIDA).

Noad T, Birnie A. 1989. *Trees of Kenya*. General Printers, Nairobi.

Robert E and Thomas T. 1999. *Challenging tradition: well-intentioned laws in Tanzania are hampering age-old practices*. *Agroforestry Today*. 11(1-2): 8-10.

**SUGGESTED CITATION**

Orwa C, A Mutua, Kindt R, Jamnadass R, S Anthony. 2009 *Agroforestry Database: a tree reference and selection guide version 4.0* (<http://www.worldagroforestry.org/sites/treedbs/treedatabases.asp>)