

Hopea odorata

white thingan, thingan, merawan

Roxb.

Dipterocarpaceae

LOCAL NAMES

Burmese (thingan net,sauchi); English (white thingan); Lao (Sino-Tibetan) (kh'e:n); Malay (merawan siput jantan,chengal pasir,chengal mas); Thai (takhian-yai,takhian-thong); Trade name (white thingan,thingan,merawan); Vietnamese (sao den,koki mosau)

BOTANIC DESCRIPTION

Hopea odorata is a medium-sized to large evergreen tree with a large crown growing to 45 m tall, bole straight, cylindrical, branchless to 25 m, with diameter of up to 4.5 m or more and prominent buttresses, bark surface scaly, grey to dark brown, longitudinally furrowed, yellow or reddish inside.

Leaves ovate-lanceolate, 7-14 by 3-7 cm, falcate, base broadly cuneate, venation scalariform, midrib applanate to slightly channeled above, glabrous on both surfaces, petiole 2 cm long, slender.

Flowers small, sweet scented, yellowish-white, very shortly pedicelled, in one-sided racemes, stamens 15, anthers narrowly ellipsoid, ovary ovoid, punctate or glabrous.

Fruit small, ovoid, wings oblanceolate, rounded, 3-4 cm long, finely veined lengthwise.

The specific epithet means odour and refers to the sweet smell of the flowers.

BIOLOGY

Hopea flowers and fruits almost regularly every two years. It is pollinated by thrips (Thysanoptera). The period between anthesis and maturity of the fruit is about three months. The small white and fragrant flowers appear between February and April and the fruits ripen at the beginning of the rainy season in May and June. The fruits are dispersed by wind and seeds germinate readily on falling to the ground.



Hopea odorata (Chongrak Wachrinrat)

Hopea odorata

Roxb.

Dipterocarpaceae

white thingan, thingan, merawan

ECOLOGY

H. odorata is a riparian species usually occurring on deep rich soils, most commonly along the banks of streams and in damp situations up to 600-m altitude. It is chiefly found in the Andamans, in moist tropical evergreen forests and occurs sporadically in pure groups, but is not gregarious over large areas. In Myanmar, it occurs in moist tropical forests.

BIOPHYSICAL LIMITS

Altitude: 0-600 m

Mean annual temperature: 25-27 deg. C

Mean annual rainfall: 1 500-3 000 mm

Soil type: It is found typically on deep rich soil, usually along the banks of streams and in damp situations.

DOCUMENTED SPECIES DISTRIBUTION

Native: Bangladesh, Cambodia, China, India, Laos, Malaysia, Myanmar, Thailand, Vietnam

Exotic:



The map above shows countries where the species has been planted. It does neither suggest that the species can be planted in every ecological zone within that country, nor that the species can not be planted in other countries than those depicted. Since some tree species are invasive, you need to follow biosafety procedures that apply to your planting site.

Hopea odorata

white thingan, thingan, merawan

Roxb.

Dipterocarpaceae

PRODUCTS

Timber: The sapwood is pale yellow or greyish yellow turning pale brown on exposure, heartwood yellowish-brown to brownish red sometimes with dark streaks, turning purplish on exposure, with lustrous white resin canals at irregular intervals, becoming dull with age. The wood is very hard and heavy weighing 755-kg/cu m, difficult to saw but finishes well. It is chiefly used for boat-building, dug-out canoes and for construction purposes, where durability and strength are of primary importance. It is also used for carts, presses flooring, roofing, piles, fence-posts, ploughs, furniture, etc. It is a first class sleeper wood.

Gum or resin: The tree yields a resin known as rock dammar in commerce, which the Burmese use to caulk boats, in painting pictures and in preparation of varnishes. A composition prepared by mixing the resin with bees-wax and red ochre is used for fastening spear and arrowheads.

Tannin or dyestuff: The leaves, bark and wood contain 11, 13-15, and 10% tannin respectively, and are used for tanning.

Medicine: The dammar is applied on sores and wounds. In Indo-China, the bark is used as a masticatory.

Other products: The bark yields a supple pale leather. Leaves have a softening effect and are used for finishing mangrove-tanned leathers.

SERVICES

Shade or shelter: The tree is sometimes used to provide shade.

Reclamation: The species is used for reforestation in Southeast Asia.

Hopea odorata

white thingan, thingan, merawan

Roxb.

Dipterocarpaceae

TREE MANAGEMENT

Germination rate has been found to be 73% in unshaded beds, 83% in shaded beds and 40% in direct field sowings. Generally, a shade crop is grown to protect the young seedlings from the first hot weather and to keep weeds down. The young plants need full overhead light and the shade crop should be cut back periodically to ensure it does not overtop the tree. Weeding, occasional watering and loosening of the soil around the plants is recommended in the nursery. Dibbling is done at an espacement of 7.5 by 7.5 cm. About 70g of seed is enough for a sq. m of nursery bed.

GERMPLASM MANAGEMENT

There are about 3 000-4 600 seeds/kg. Seeds are recalcitrant and die within five days due to dehydration. If dried at 35 deg. C to 33 % moisture content, seeds could stay viable for 1-2 months at 15 deg. C, maintaining a germination rate of over 60 %. If stored at 4 deg. C, the seeds can stay viable for about three months.

PESTS AND DISEASES

Attacks by defoliators have frequently been noticed in plantations towards the close of the rains and continue until the end of the hot weather when new and healthy leaves appear. Stem borers attack saplings in the natural forest. The weevil *Nanophyes shorea* attacks seeds. Several beetles and larvae of insects of the orders Coleoptera and Isoptera bore in the dead wood and fallen wood.

Hopea odorata

white thingan, thingan, merawan

Roxb.

Dipterocarpaceae

FURTHER READING

Oldfields S, Lusty C, Mackinnen A. 1998. The world list of threatened trees. 650 pp. World Conservation Press, Cambridge, UK.

Soerianegara I, Lemmens RHMJ (eds.). 1993. Plant Resources of South-East Asia. No. 5(1): Timber trees: major commercial timbers. Backhuys Publishers, Leiden.

Tixier P. 1973. The biology of the Dipterocarpaceae: phenology and germination of *Hopea odorata* at Dangkor (Kandal). Bois et Forêts des Tropiques. 48: 47-52.

Troup RS. 1929. The silviculture of Indian trees. Controller of publications, New Delhi, India.

Troup RS. 1975. The silviculture of Indian trees. ed. 2, vol. 1. Government of India.

SUGGESTED CITATION

Orwa C, A Mutua, Kindt R, Jamnadass R, S Anthony. 2009 Agroforestry Database: a tree reference and selection guide version 4.0 (<http://www.worldagroforestry.org/sites/treedbs/treedatabases.asp>)